



Montgomery County

Tobacco/ESD Use by Adults: 2012-2017

Behavioral Risk Factor Surveillance System (BRFSS)

CURRENT USE OF TOBACCO/ELECTRONIC SMOKING DEVICES (ESDs)				
<i>Estimated Prevalence (%)</i>	2012	2014	2016	2017*
<i>Confidence Interval (CI)</i>	% CI N	% CI N	% CI N	% CI N
<i>Estimated Number (N)</i>				
Any Tobacco <i>(Any tobacco includes cigarettes, cigars, smokeless tobacco, and other tobacco products, excludes ESDs)</i>	10.9 (8.2-13.7) 81,828	10.3 (7.6-13.0) 82,171	8.4 (6.7-10.1) 68,320	9.5 (7.3-11.7) 77,041
Cigarettes	7.5 (5.5-9.6) 55,857	7.9 (5.3-10.5) 59,938	6.2 (4.7-7.8) 47,185	7.3 (5.3-9.3) 56,338
Cigars	BRFSS Data Not Available	2.9 (1.5-4.4) 20,235	1.5 (0.7-2.3) 10,217	3.5 (1.8-5.2) 23,613
Smokeless Tobacco	BRFSS Data Not Available	BRFSS Data Not Available	1.1 (0.5-1.8) 8,783	BRFSS Data Not Available
ESDs	No BRFSS Data Collected	1.8 (0.8-2.8) 12,554	1.8 (0.9-2.6) 13,394	3.1 (1.8-4.5) 23,642
CURRENT TOBACCO USE — Gender and Race/Ethnicity				
<i>Estimated Prevalence (%)</i>	2012	2014	2016	2017*
<i>Confidence Interval (CI)</i>	% CI N	% CI N	% CI N	% CI N
<i>Estimated Number (N)</i>				
Female	7.1 (4.4-9.8) 25,205	5.8 (3.3-8.3) 24,210	4.3 (2.9-5.6) 18,249	7.9 (5.5-10.3) 33,857
Male	14.3 (10.1-18.5) 63,964	15.4 (10.5-20.3) 57,960	13.0 (9.8-16.2) 50,071	11.2 (7.4-14.9) 43,184
White	13.9 (10.1-17.7) 53,149	9.5 (6.7-12.2) 36,477	8.0 (5.7-10.4) 30,226	8.7 (6.1-11.3) 32,281
African American/Black	BRFSS Data Not Available	BRFSS Data Not Available	9.8 (5.8-13.9) 13,647	17.2 (9.8-24.6) 24,479
Asian	BRFSS Data Not Available	BRFSS Data Not Available	BRFSS Data Not Available	BRFSS Data Not Available
Hispanic/Latino	BRFSS Data Not Available	BRFSS Data Not Available	BRFSS Data Not Available	BRFSS Data Not Available
American Indian/Alaskan Native	BRFSS Data Not Available	BRFSS Data Not Available	BRFSS Data Not Available	BRFSS Data Not Available
**Minority Combined (Race & Female)	8.0 (5.4-10.7) 47,808	9.2 (5.9-12.5) 55,714	7.2 (5.4-8.9) 45,165	8.9 (6.3-11.4) 55,694

When "BRFSS Data Not Available" is noted in the tables, this is due to a high uncertainty or an insufficient sample size.

*Any tobacco includes cigarettes, cigars, and smokeless tobacco, excludes ESDs and other tobacco (pipes, hookah, bidis, kreteks, or dissolvable).

**By CRF/Tobacco legislative statute, minority is defined as an individual of African American, Hispanic, Native American, or Asian descent plus all women (MD Health-General Article §13-1101).

CURRENT TOBACCO USE — Income and Education					
<i>Estimated Prevalence (%)</i>	2012	2014	2016	2017*	
<i>Confidence Interval (CI)</i>	% CI N	% CI N	% CI N	% CI N	
<i>Estimated Number (N)</i>					
Income					
< \$50k	11.3 (6.3-16.3) 26,520	10.8 (5.3-16.2) 22,007	12.6 (8.0-17.3) 24,771	12.3 (7.2-17.3) 26,251	
> \$50k	11.7 (8.3-15.1) 52,884	10.9 (7.5-14.2) 46,975	7.0 (5.2-8.9) 33,137	8.5 (5.7-11.3) 39,339	
Education					
No High School	BRFSS Data Not Available	BRFSS Data Not Available	BRFSS Data Not Available	BRFSS Data Not Available	
High School or GED	15.2 (7.4-23.1) 20,640	16.2 (8.2-24.3) 21,874	12.1 (6.6-17.6) 13,217	24.2 (15.5-32.9) 31,753	
Some College	15.9 (9.3-22.4) 26,451	12.6 (6.3-19.0) 21,509	12.3 (7.6-16.9) 21,891	10.4 (5.3-15.5) 17,439	
College Grad	7.4 (5.0-9.9) 29,684	6.8 (4.7-8.9) 26,786	5.0 (3.5-6.5) 21,723	4.5 (3.0-5.9) 18,835	
HOUSEHOLDS WITH MINOR CHILDREN THAT HAVE RESIDENT ADULT SMOKERS					
<i>Estimated Prevalence (%)</i>	2012	2014	2016	2017	
<i>Confidence Interval (CI)</i>	% CI N	% CI N	% CI N	% N	
<i>Estimated Number (N)</i>					
Proportion with Adult Smokers	15.2 (9.7-20.8) 41,554	11.3 (6.5-16.2) 31,651	8.5 (5.4-11.6) 22,479	BRFSS Data Not Collected	
CIGARETTE SMOKING AMONG THOSE RECEIVING MENTAL HEALTH AND/OR SUBSTANCE-RELATED DISORDER SERVICES					
Maryland Outcomes Measurement System Datamart (OMS)					
<i>Prevalence (%)</i>	2014	2015	2016	2017	2018
<i>Number (N)</i>	% N	% N	% N	% N	% N
Mental Health Services	33.7 (1,291)	28.8 (1,234)	28.6 (1,348)	24.2 (1,187)	20.0 (1,039)
Substance-Related Disorder Services	Not Available	55.2 (148)	62.1 (308)	65.0 (575)	59.0 (563)
Both	Not Available	62.4 (58)	68.4 (121)	67.0 (203)	65.0 (160)
PERCENT AND NUMBER OF PREGNANT FEMALES SMOKING DURING PREGNANCY					
Vital Statistics Administration					
<i>Prevalence (%)</i>	2012	2014	2016	2017	
<i>Number (N)</i>	% N	% N	% N	% N	
Proportion of Pregnant Smokers	1.6 (213)	1.4 (186)	1.0 (134)	1.0 (132)	

When "BRFSS Data Not Available" is noted in the tables, this is due to a high uncertainty or an insufficient sample size.

*Any tobacco includes cigarettes, cigars, and smokeless tobacco, excludes ESDs and other tobacco (pipes, hookah, bidis, kreteks, or dissolvable).

**By CRF/Tobacco legislative statute, minority is defined as an individual of African American, Hispanic, Native American, or Asian descent plus all women (MD Health-General Article §13-1101).