



Prince George's County

Tobacco/ESD Use by Adults: 2012-2017

Behavioral Risk Factor Surveillance System (BRFSS)

CURRENT USE OF TOBACCO/ELECTRONIC SMOKING DEVICES (ESDs)				
<i>Estimated Prevalence (%)</i>	2012	2014	2016	2017*
<i>Confidence Interval (CI)</i>	% CI N	% CI N	% CI N	% CI N
<i>Estimated Number (N)</i>				
Any Tobacco <i>(Any tobacco includes cigarettes, cigars, smokeless tobacco, and other tobacco products, excludes ESDs)</i>	16.6 (13.1-20.1) 103,805	18.3 (14.1-22.4) 127,102	13.6 (11.3-15.9) 97,260	13.4 (10.7-16.1) 95,279
Cigarettes	14.2 (10.8-17.5) 88,296	11.8 (8.7-15.0) 79,628	11.3 (9.1-13.4) 74,789	10.3 (7.8-12.7) 68,564
Cigars	4.7 (2.6-6.7) 23,783	8.1 (4.5-11.6) 52,011	3.1 (2.0-4.3) 18,610	4.1 (2.5-5.6) 25,441
Smokeless Tobacco	BRFSS Data Not Available	BRFSS Data Not Available	BRFSS Data Not Available	BRFSS Data Not Available
ESDs	No BRFSS Data Collected	BRFSS Data Not Available	2.4 (1.3-3.5) 15,768	BRFSS Data Not Available
CURRENT TOBACCO USE — Gender and Race/Ethnicity				
<i>Estimated Prevalence (%)</i>	2012	2014	2016	2017*
<i>Confidence Interval (CI)</i>	% CI N	% CI N	% CI N	% CI N
<i>Estimated Number (N)</i>				
Female	13.1 (9.5-16.8) 43,722	13.6 (9.2-18.0) 49,837	10.0 (7.4-12.6) 37,598	9.3 (6.4-12.1) 34,659
Male	21.1 (15.0-27.1) 64,038	23.5 (16.4-30.7) 77,266	17.7 (13.8-21.5) 59,662	18.0 (13.3-22.7) 60,619
White	18.5 (11.5-25.5) 23,520	16.7 (9.3-24.2) 19,367	14.6 (9.7-19.4) 16,394	14.2 (9.5-18.9) 15,567
African American/Black	16.9 (12.8-20.9) 66,271	19.9 (14.4-25.5) 85,302	15.0 (11.9-18.0) 65,044	12.6 (9.3-16.0) 54,833
Asian	BRFSS Data Not Available	BRFSS Data Not Available	BRFSS Data Not Available	BRFSS Data Not Available
Hispanic/Latino	BRFSS Data Not Available	BRFSS Data Not Available	BRFSS Data Not Available	16.5 (7.2-25.8) 18,951
American Indian/Alaskan Native	BRFSS Data Not Available	BRFSS Data Not Available	BRFSS Data Not Available	BRFSS Data Not Available
**Minority Combined (Race & Female)	16.4 (12.8-20.1) 93,972	17.7 (13.3-22.1) 113,264	13.1 (10.7-15.4) 85,814	13.0 (10.2-15.9) 85,325

When "BRFSS Data Not Available" is noted in the tables, this is due to a high uncertainty or an insufficient sample size.

*Any tobacco includes cigarettes, cigars, and smokeless tobacco, excludes ESDs and other tobacco (pipes, hookah, bidis, kreteks, or dissolvable).

**By CRF/Tobacco legislative statute, minority is defined as an individual of African American, Hispanic, Native American, or Asian descent plus all women (MD Health-General Article §13-1101).

CURRENT TOBACCO USE — Income and Education				
<i>Estimated Prevalence (%)</i>	2012	2014	2016	2017*
<i>Confidence Interval (CI)</i>	% CI N	% CI N	% CI N	% CI N
<i>Estimated Number (N)</i>				
Income				
< \$50k	23.3 (16.5-30.1)	17.4 (10.5-24.3)	16.0 (11.9-20.1)	17.7 (11.9-23.5)
	56,923	42,110	37,900	42,463
> \$50k	12.3 (8.9-15.7)	19.0 (13.1-25.0)	13.2 (9.9-16.5)	13.5 (9.8-17.2)
	38,252	63,513	43,266	45,298
Education				
No High School	28.7 (13.1-44.3)	BRFSS Data Not Available	BRFSS Data Not Available	BRFSS Data Not Available
	22,374			
High School or GED	24.8 (17.0-32.5)	24.3 (15.2-33.4)	19.0 (14.1-23.9)	18.7 (12.5-24.8)
	39,769	43,416	34,392	34,284
Some College	15.8 (10.0-21.6)	19.5 (12.6-26.5)	13.8 (9.8-17.8)	13.0 (8.2-17.9)
	32,952	46,655	32,436	28,302
College Grad	6.8 (4.3-9.2)	7.4 (4.1-10.7)	7.8 (5.3-10.3)	9.8 (6.2-13.4)
	12,664	14,546	16,157	22,216
HOUSEHOLDS WITH MINOR CHILDREN THAT HAVE RESIDENT ADULT SMOKERS				
<i>Estimated Prevalence (%)</i>	2012	2014	2016	2017
<i>Confidence Interval (CI)</i>	% CI N	% CI N	% CI N	% CI N
<i>Estimated Number (N)</i>				
Proportion with Adult Smokers	21.4 (14.7-28.2)	17.4 (11.3-23.5)	20.1 (14.8-25.5)	BRFSS Data Not Collected
	48,264	44,688	45,308	
CIGARETTE SMOKING AMONG THOSE RECEIVING MENTAL HEALTH AND/OR SUBSTANCE-RELATED DISORDER SERVICES				
Maryland Outcomes Measurement System Datamart (OMS)				
<i>Prevalence (%)</i>	2014	2015	2016	2017
<i>Number (N)</i>	% N	% N	% N	% N
Mental Health Services	32.9 (1,285)	29.3 (1,371)	26.1 (1,298)	22.8 (1,225)
	19.2 (1,133)			
Substance-Related Disorder Services	Not Available	46.3 (202)	57.1 (382)	53.9 (578)
		54.2 (567)		
Both	Not Available	56.5 (52)	53.1 (69)	56.4 (119)
		50.6 (90)		
PERCENT AND NUMBER OF PREGNANT FEMALES SMOKING DURING PREGNANCY				
Vital Statistics Administration				
<i>Prevalence (%)</i>	2012	2014	2016	2017
<i>Number (N)</i>	% N	% N	% N	% N
Proportion of Pregnant Smokers	2.4 (281)	2.5 (304)	1.6 (197)	1.6 (200)

When "BRFSS Data Not Available" is noted in the tables, this is due to a high uncertainty or an insufficient sample size.

*Any tobacco includes cigarettes, cigars, and smokeless tobacco, excludes ESDs and other tobacco (pipes, hookah, bidis, kreteks, or dissolvable).

**By CRF/Tobacco legislative statute, minority is defined as an individual of African American, Hispanic, Native American, or Asian descent plus all women (MD Health-General Article §13-1101).