



# Washington County

## Tobacco/ESD Use by Adults: 2012-2017

Behavioral Risk Factor Surveillance System (BRFSS)

<b>CURRENT USE OF TOBACCO/ELECTRONIC SMOKING DEVICES (ESDs)</b>				
<i>Estimated Prevalence (%)</i>	<b>2012</b>	<b>2014</b>	<b>2016</b>	<b>2017*</b>
<i>Confidence Interval (CI)</i>	<b>% CI N</b>	<b>% CI N</b>	<b>% CI N</b>	<b>% CI N</b>
<i>Estimated Number (N)</i>				
<b>Any Tobacco</b> <i>(Any tobacco includes cigarettes, cigars, smokeless tobacco, and other tobacco products, excludes ESDs)</i>	<b>20.8</b> (14.0-27.6) <b>21,800</b>	<b>24.7</b> (16.9-32.5) <b>29,014</b>	<b>21.9</b> (17.5-26.4) <b>25,656</b>	<b>19.6</b> (13.8-25.4) <b>23,120</b>
<b>Cigarettes</b>	<b>18.0</b> (11.3-24.7) <b>18,809</b>	<b>21.5</b> (14.0-29.1) <b>24,533</b>	<b>18.3</b> (14.2-22.4) <b>20,347</b>	<b>17.3</b> (11.6-23.0) <b>19,461</b>
<b>Cigars</b>	BRFSS Data Not Available	BRFSS Data Not Available	BRFSS Data Not Available	BRFSS Data Not Available
<b>Smokeless Tobacco</b>	BRFSS Data Not Available	BRFSS Data Not Available	BRFSS Data Not Available	BRFSS Data Not Available
<b>ESDs</b>	No BRFSS Data Collected	BRFSS Data Not Available	<b>4.2</b> (2.0-6.5) <b>4,647</b>	<b>4.2</b> (1.9-6.4) <b>4,594</b>
<b>CURRENT TOBACCO USE — Gender and Race/Ethnicity</b>				
<i>Estimated Prevalence (%)</i>	<b>2012</b>	<b>2014</b>	<b>2016</b>	<b>2017*</b>
<i>Confidence Interval (CI)</i>	<b>% CI N</b>	<b>% CI N</b>	<b>% CI N</b>	<b>% CI N</b>
<i>Estimated Number (N)</i>				
<b>Female</b>	<b>16.3</b> (11.0-21.6) <b>9,526</b>	<b>19.6</b> (9.8-29.3) <b>11,226</b>	<b>19.4</b> (14.1-24.7) <b>11,183</b>	<b>16.8</b> (9.8-23.8) <b>9,791</b>
<b>Male</b>	<b>27.1</b> (14.6-39.7) <b>13,386</b>	<b>29.6</b> (17.6-41.7) <b>17,788</b>	<b>24.4</b> (17.3-31.4) <b>14,473</b>	<b>22.3</b> (13.1-31.6) <b>13,330</b>
<b>White</b>	<b>18.5</b> (12.8-24.2) <b>17,444</b>	<b>25.1</b> (16.7-33.5) <b>23,948</b>	<b>22.4</b> (17.4-27.3) <b>21,617</b>	<b>21.4</b> (14.8-27.9) <b>21,489</b>
<b>African American/Black</b>	BRFSS Data Not Available	BRFSS Data Not Available	BRFSS Data Not Available	BRFSS Data Not Available
<b>Asian</b>	BRFSS Data Not Available	BRFSS Data Not Available	BRFSS Data Not Available	BRFSS Data Not Available
<b>Hispanic/Latino</b>	BRFSS Data Not Available	BRFSS Data Not Available	BRFSS Data Not Available	BRFSS Data Not Available
<b>American Indian/Alaskan Native</b>	BRFSS Data Not Available	BRFSS Data Not Available	BRFSS Data Not Available	BRFSS Data Not Available
<b>**Minority Combined (Race &amp; Female)</b>	<b>20.8</b> (12.3-29.4) <b>13,617</b>	<b>22.0</b> (12.3-31.8) <b>15,884</b>	<b>19.2</b> (14.1-24.2) <b>12,853</b>	<b>16.2</b> (9.8-22.6) <b>11,174</b>

When "BRFSS Data Not Available" is noted in the tables, this is due to a high uncertainty or an insufficient sample size.

\***Any tobacco** includes cigarettes, cigars, and smokeless tobacco, excludes ESDs and other tobacco (pipes, hookah, bidis, kreteks, or dissolvable).

\*\*By CRF/Tobacco legislative statute, minority is defined as an individual of African American, Hispanic, Native American, or Asian descent plus all women (MD Health-General Article §13-1101).

<b>CURRENT TOBACCO USE — Income and Education</b>				
<i>Estimated Prevalence (%)</i>	<b>2012</b>	<b>2014</b>	<b>2016</b>	<b>2017*</b>
<i>Confidence Interval (CI)</i>	<b>% CI N</b>	<b>% CI N</b>	<b>% CI N</b>	<b>% CI N</b>
<i>Estimated Number (N)</i>				
<b>Income</b>				
<b>&lt; \$50k</b>	<b>33.2</b> (19.9-46.4)	<b>28.2</b> (16.8-39.5)	<b>31.2</b> (23.4-38.9)	<b>23.8</b> (13.0-34.7)
	<b>14,921</b>	<b>18,301</b>	<b>14,634</b>	<b>11,239</b>
<b>&gt; \$50k</b>	<b>10.1</b> (5.1-15.1)	<b>23.4</b> (10.8-36.0)	<b>16.0</b> (9.6-22.3)	<b>19.5</b> (10.5-28.4)
	<b>4,989</b>	<b>8,772</b>	<b>7,853</b>	<b>8,951</b>
<b>Education</b>				
<b>No High School/High School- GED</b>	<b>27.1</b> (16.7-37.6)	<b>33.8</b> (21.6-45.9)	<b>28.8</b> (21.8-35.7)	<b>22.3</b> (13.8-30.8)
	<b>16,748</b>	<b>20,127</b>	<b>18,212</b>	<b>15,276</b>
<b>Some College/College Grad</b>	<b>12.4</b> (6.9-17.8)	<b>15.4</b> (6.8-24.0)	<b>13.9</b> (8.9-18.9)	<b>15.9</b> (8.5-23.3)
	<b>5,607</b>	<b>8,887</b>	<b>7,444</b>	<b>7,845</b>
<b>HOUSEHOLDS WITH MINOR CHILDREN THAT HAVE RESIDENT ADULT SMOKERS</b>				
<i>Estimated Prevalence (%)</i>	<b>2012</b>	<b>2014</b>	<b>2016</b>	<b>2017</b>
<i>Confidence Interval (CI)</i>	<b>% CI N</b>	<b>% CI N</b>	<b>% CI N</b>	<b>% CI N</b>
<i>Estimated Number (N)</i>				
<b>Proportion with Adult Smokers</b>	<b>30.0</b> (16.7-43.3)	<b>44.0</b> (28.5-59.5)	<b>34.8</b> (25.9-43.7)	<b>BRFSS Data Not Collected</b>
	<b>12,818</b>	<b>16,240</b>	<b>14,099</b>	
<b>CIGARETTE SMOKING AMONG THOSE RECEIVING MENTAL HEALTH AND/OR SUBSTANCE-RELATED DISORDER SERVICES</b>				
Maryland Outcomes Measurement System Datamart (OMS)				
<i>Prevalence (%)</i>	<b>2014</b>	<b>2015</b>	<b>2016</b>	<b>2017</b>
<i>Number (N)</i>	<b>% N</b>	<b>% N</b>	<b>% N</b>	<b>% N</b>
<b>Mental Health Services</b>	<b>53.4</b> (1,394)	<b>51.3</b> (1,627)	<b>51.2</b> (1,696)	<b>50.8</b> (1,759)
<b>Substance-Related Disorder Services</b>	<b>Not Available</b>	<b>73.9</b> (441)	<b>74.7</b> (892)	<b>75.7</b> (1,752)
		<b>70.1</b> (1,649)		
<b>Both</b>	<b>Not Available</b>	<b>74.2</b> (181)	<b>70.2</b> (250)	<b>77.4</b> (432)
				<b>68.0</b> (382)
<b>PERCENT AND NUMBER OF PREGNANT FEMALES SMOKING DURING PREGNANCY</b>				
Vital Statistics Administration				
<i>Prevalence (%)</i>	<b>2012</b>	<b>2014</b>	<b>2016</b>	<b>2017</b>
<i>Number (N)</i>	<b>% N</b>	<b>% N</b>	<b>% N</b>	<b>% N</b>
<b>Proportion of Pregnant Smokers</b>	<b>21.3</b> (376)	<b>20.0</b> (361)	<b>17.4</b> (294)	<b>16.1</b> (264)

When "BRFSS Data Not Available" is noted in the tables, this is due to a high uncertainty or an insufficient sample size.

\*Any tobacco includes cigarettes, cigars, and smokeless tobacco, excludes ESDs and other tobacco (pipes, hookah, bidis, kreteks, or dissolvable).

\*\*By CRF/Tobacco legislative statute, minority is defined as an individual of African American, Hispanic, Native American, or Asian descent plus all women (MD Health-General Article §13-1101).