



Allegany County

Tobacco/ESD Use by Adults: 2012-2018

Behavioral Risk Factor Surveillance System (BRFSS)

| CURRENT USE OF TOBACCO/ELECTRONIC SMOKING DEVICES (ESDs) | | | | |
|---|---|---|---|---|
| <i>Estimated Prevalence (%)</i> | 2012* | 2014 | 2016 | 2018 |
| <i>Confidence Interval (CI)</i> | % CI N | % CI N | % CI N | % CI N |
| <i>Estimated Number (N)</i> | | | | |
| Any Tobacco <i>(Any tobacco includes cigarettes, cigars, smokeless tobacco, and other tobacco products, and ESDs)</i> | 28.3 (17.9-38.7) 17,530 | 24.6 (16.2-33.1) 13,734 | 24.9 (18.5-31.3) 13,304 | 31.8 (24.2-39.5) 19,256 |
| Cigarettes | 24.4 (14.0-34.8) 14,981 | 18.7 (11.0-26.3) 10,301 | 15.6 (10.0-21.1) 8,166 | 20.5 (13.1-27.9) 11,688 |
| Cigars | BRFSS Data Not Available | BRFSS Data Not Available | BRFSS Data Not Available | BRFSS Data Not Available |
| Smokeless Tobacco | BRFSS Data Not Available | BRFSS Data Not Available | BRFSS Data Not Available | 6.0 (2.8-9.3) 3,432 |
| ESDs | No BRFSS Data Collected | BRFSS Data Not Available | BRFSS Data Not Available | 9.4 (4.3-14.6) 5,316 |
| CURRENT TOBACCO USE — Gender and Race/Ethnicity | | | | |
| <i>Estimated Prevalence (%)</i> | 2012* | 2014 | 2016 | 2018 |
| <i>Confidence Interval (CI)</i> | % CI N | % CI N | % CI N | % CI N |
| <i>Estimated Number (N)</i> | | | | |
| Female | 24.6 (11.8-37.5) 7,716 | 20.1 (11.0-29.1) 6,173 | 15.9 (9.9-21.8) 4,457 | 31.3 (20.6-41.9) 10,464 |
| Male | 30.5 (14.8-46.2) 9,948 | 30.3 (15.7-44.9) 7,561 | 35.0 (24.0-46.0) 8,847 | 32.6 (21.6-43.6) 8,791 |
| White | 29.1 (18.2-39.9) 16,514 | 26.3 (17.3-35.4) 13,099 | 22.7 (16.3-29.1) 10,643 | 31.9 (23.9-39.8) 18,270 |
| African American/Black | BRFSS Data Not Available | BRFSS Data Not Available | BRFSS Data Not Available | BRFSS Data Not Available |
| Asian | BRFSS Data Not Available | BRFSS Data Not Available | BRFSS Data Not Available | BRFSS Data Not Available |
| Hispanic/Latino | BRFSS Data Not Available | BRFSS Data Not Available | BRFSS Data Not Available | BRFSS Data Not Available |
| American Indian/Alaskan Native | BRFSS Data Not Available | BRFSS Data Not Available | BRFSS Data Not Available | BRFSS Data Not Available |
| **Minority Combined (Race & Female) | 21.9 (10.2-33.7) 8,047 | 20.1 (11.3-29.0) 6,430 | 20.4 (13.0-27.7) 6,338 | 31.0 (20.5-41.5) 10,579 |

When "BRFSS Data Not Available" is noted in the tables, this is due to a high uncertainty or an insufficient sample size.

*Data on ESDs not collected during the survey year.

**By CRF/Tobacco legislative statute, minority is defined as an individual of African American, Hispanic, Native American, or Asian descent plus all women (MD Health-General Article §13-1101).

| CURRENT TOBACCO USE — Income and Education | | | | |
|--|---|--|--|---|
| <i>Estimated Prevalence (%)</i> | 2012* | 2014 | 2016 | 2018 |
| <i>Confidence Interval (CI)</i> | % CI N | % CI N | % CI N | % CI N |
| <i>Estimated Number (N)</i> | | | | |
| Income | | | | |
| < \$50k | 34.7 (20.4-49.0) 12,050 | 35.2 (21.9-48.5) 9,616 | 25.3 (16.3-34.4) 6,841 | 36.5 (24.6-48.4) 11,041 |
| > \$50k | BRFSS Data Not Available | BRFSS Data Not Available | 18.8 (9.6-28.1) 2,510 | 31.1 (19.4-42.8) 6,758 |
| Education | | | | |
| No High School/High School- GED | 22.0 (11.6-32.5) 7,718 | 26.3 (14.1-38.5) 8,256 | 23.3 (14.8-31.8) 5,597 | 44.5 (32.3-56.7) 12,409 |
| Some College/College Grad | 34.5 (17.4-51.6) 9,946 | 22.6 (11.3-33.9) 5,478 | 26.3 (16.9-35.6) 7,707 | 21.2 (13.5-28.9) 6,846 |
| HOUSEHOLDS WITH MINOR CHILDREN THAT HAVE RESIDENT ADULT SMOKERS | | | | |
| <i>Estimated Prevalence (%)</i> | 2012 | 2014 | 2016 | 2018 |
| <i>Confidence Interval (CI)</i> | % CI N | % CI N | % CI N | % CI N |
| <i>Estimated Number (N)</i> | | | | |
| Proportion with Adult Smokers | 31.5 (14.2-48.8) 4,496 | 44.6 (23.8-65.3) 7,256 | 40.1 (23.6-56.7) 5,446 | 39.9 (21.0-58.8) 6,282 |
| CIGARETTE SMOKING AMONG THOSE RECEIVING MENTAL HEALTH AND/OR SUBSTANCE USE DISORDER SERVICES (CURRENT AS OF 4/2019) | | | | |
| Maryland Outcomes Measurement System Datamart (OMS) | | | | |
| <i>Prevalence (%)</i> | 2014 | 2015 | 2016 | 2017 |
| <i>Number (N)</i> | % N | % N | % N | % N |
| Mental Health Services | 55.2 (790) | 53.1 (736) | 54.9 (757) | 42.8 (605) |
| Substance Use Disorder Services | Not Available | 77.2 (287) | 79.4 (676) | 77.0 (1,037) |
| Both | Not Available | 79.6 (86) | 79.4 (154) | 65.3 (192) |
| | | | | 59.1 (176) |
| PERCENT AND NUMBER OF PREGNANT FEMALES SMOKING DURING PREGNANCY | | | | |
| Vital Statistics Administration | | | | |
| <i>Prevalence (%)</i> | 2012 | 2014 | 2016 | 2018 |
| <i>Number (N)</i> | % N | % N | % N | % N |
| Proportion of Pregnant Smokers | 26.7 (186) | 24.3 (161) | 24.7 (153) | 23.5 (158) |

When "BRFSS Data Not Available" is noted in the tables, this is due to a high uncertainty or an insufficient sample size.

*Data on ESDs not collected during the survey year.

**By CRF/Tobacco legislative statute, minority is defined as an individual of African American, Hispanic, Native American, or Asian descent plus all women (MD Health-General Article §13-1101).